



Evaluation of the prevalence of daytime lethargy, circadian inclination, caffeine utilization and utilization of different energizers among Thai understudies

Takougang I.¹, Ngogang J.¹ and Sihom F.²

¹Department of Epidemiology, Multidisciplinary Health International Research Training Program, Harvard School of Public Health, Boston, MA, U.S.A.

²Department of Mathematics and Statistics, Georgetown University, Washington, DC, U.S.A.

Abstract

This study was conducted to evaluate the prevalence of daytime sleepiness and evening chronotype and to assess the extent to which both are associated with the use of caffeinated stimulants among 3,000 Thai college students. Demographic and behavioral characteristics were collected using a self-administered questionnaire. The Epworth Sleepiness Scale and the Horne and Ostberg Morningness-Eveningness Questionnaire were used to evaluate prevalence of daytime sleepiness and circadian preference. Multivariable logistic regression models were used to evaluate the association between sleep habits and consumption of caffeinated beverages. Overall, the prevalence of daytime sleepiness was 27.9% (95% CI: 26.2 to 29.5%) while the prevalence of evening chronotype was 13.0% (95% CI: 11.8 to 14.2%). Students who use energy drinks were more likely to be evening types. For instance, the use of M100/M150 energy drinks was associated with a more than 3-fold increased odds of evening chronotype (OR 3.50; 95% CI 1.90 to 6.44), while Red Bull users were more than twice as likely to have evening chronotype (OR 2.39; 95% CI 1.02 to 5.58). Additionally, those who consumed any energy drinks were more likely to be daytime sleepers. For example, Red Bull (OR 1.72; 95% CI 1.08 to 2.75) or M100/M150 (OR 1.52; 95% CI 1.10 to 2.11) consumption was associated with increased odds of daytime sleepiness. Our findings emphasize the importance of implementing educational and prevention programs targeted toward improving sleep hygiene and reducing the consumption of energy drinks among young adults.

Keywords: Daytime sleepiness, Caffeine, stimulants, college students, morningness-eveningness.

INTRODUCTION

Sleep is essential for maintaining good health in humans (Luyster et al., 2012). However, adolescents experience

many changes in their sleep patterns in part due to physiological delayed sleep phase and disruption of the homeostatic sleep-wake cycle (Millman, 2005; Oginska and Pokorski, 2006). The challenges associated with the transition to college, including reduced parental supervision, extracurricular activities, academic workloads and social commitments (Malinauskas et al., 2007; Millman, 2005; Oginska et al., 2006) adversely affect sleep duration and sleep patterns of college students (Lund et al., 2010; Millman, 2005). Irregular sleep patterns can lead to daytime sleepiness, which is a result of sleep loss and deprivation (AASM, 2001; Kushida, 2006). There is a substantial body of evidence that shows daytime sleepiness is associated with reduced emotional intelligence, impaired constructive thinking skills, poor academic performance, job loss, headaches and obesity (AASM, 2001; Killgore et al., 2008; Slater et al., 2013).

Additionally, circadian rhythms and preferences can also be affected by social, biological and environmental factors (Besoluk et al., 2011; Kanerva et al., 2012; Lucassen et al., 2013). There are two major types of circadian preferences: morningness and eveningness (Horne and Ostberg, 1976; Schneider et al., 2011). Morning types tend to wake up relatively early in the morning and work best during this time (Schneider et al., 2011). On the other hand, evening types wake up later in the day and prefer to work during the evening and night (Schneider et al., 2011). An accumulating body of literature has confirmed that evening types experience higher sleep apnea, increased stress hormones, less healthier lifestyles and lower academic performances, although much of the research has focused on Western populations (Besoluk et al., 2011; Kanerva et al., 2012; Lucassen et al., 2013; Merikanto et al., 2013).

Caffeinated beverages and energy drinks have been implicated as important risk factors for increased daytime sleepiness and evening chronotype among college students (Ishak et al., 2012; Taillard et al., 1999). Globally, energy drinks have gained popularity among adolescents and young adults to counteract tiredness and meet academic, physical and cognitive demands (Buxton and Hagan, 2012; Malinauskas et al., 2007; Pomeranz et al., 2013; Roehrs and Roth, 2008). Some adverse effects of caffeine intake include energy loss, headaches, cardiac problems and even sudden death (Cannon et al., 2001; Roehrs and Roth, 2008; Seifert et al., 2011).

Despite the rising trends and aggressive marketing strategies aimed toward college students, there has been little research done on college students' intake of energy drinks and their impact on sleep disorders (Arria et al., 2010; Malinauskas et al., 2007). In particular, Thailand leads in consumption of energy drink consumption per capita globally, but very few studies have focused on sleep-related health problems in Southeast Asian populations (Cheng et al., 2012; Doi et al., 2003; Mak et al., 2010; Tsai and Li, 2004; Zenith International, 2007).

Given documented relationships between energy drinks

and sleep disorders, we hypothesized that students who use more stimulants were more likely than non-users to experience daytime sleepiness (Ishak et al., 2012; James et al., 2011; Roehrs and Roth 2008). We also hypothesized that students who use stimulants are more likely than non-users to be evening types (Taillard et al., 1999). From this study, we expect to provide evidence that will guide the development of health and wellness programs for young adults in Thailand.

MATERIALS AND METHODS

Study setting and sample

This cross-sectional study was conducted between December, 2010 and February, 2011 at seven private and government colleges in Thailand (Lohsoonthorn et al., 2012). Details of the study setting, sampling and data collection procedures have been described previously (Lohsoonthorn et al., 2012). Briefly, flyers were posted in each campus to recruit and invite participants to the study. Students who expressed an interest in participating were asked to meet in a large classroom or an auditorium where they were informed about the purpose of the study. Students consenting to participate were asked to complete a self-administered individual survey. Vision impaired students and those who could not read the consent and questionnaire forms were not eligible to participate. Those enrolled in correspondence, extension or night school programs were not included as well since their experience might be different from regular time students. Approximately 90% students invited to participate in the survey elected to do so. A total of 3,000 undergraduate students participated in the study and completed a self-administered survey. All study questionnaires were anonymous, and no personal identifiers were collected. Study procedures were approved by the institutional review boards of the Faculty of Medicine Chulalongkorn University, Walailak University and the University of Washington, USA. The Harvard School of Public Health Office of Human Research Administration, USA, granted approval to use the anonymous data set for analysis.

Data collection and covariates

Demographics

In this study, a self-administered questionnaire was used to collect information on demographics and lifestyle characteristics. These include age, sex, education level, smoking status, physical activity, caffeine use, and alcohol consumption. The student's height, weight, waist and hip circumference were measured by research nurses after the questionnaire was administered.

Use of stimulant beverages and other caffeinated drinks

Participants were first asked if they consumed any stimulant or energy drink during the past week. Participants answering "Yes" were further asked to identify the specific types of energy drinks and/or stimulant drinks. These are beverages used to provide an extra boost in energy, promote wakefulness and provide cognitive and mood enhancement (Ishak et al., 2012). In this study, we will be using the terms energy drinks and stimulant beverages interchangeably. To provide a range of popular energy drinks in the immediate geographic regions where the survey was administered, we included examples of energy drinks that were common on the

campuses and in surrounding social establishments. These included global and local brands such as Red Bull, M100, M150, CarabaoDaeng, Lipovitan-D or Lipo, Wrangyer and Sharks. For the purpose of this analysis, we combined energy drinks that were less commonly used (that is, CarabaoDaeng, Lipovitan-D or Lipo, Wrangyer and Shark) and classified them as “other energy drinks”. Consumption of other caffeine-containing beverages included coffee, black tea and stimulant beverages such as Coke and Pepsi or sugar-free Coke and Pepsi (contain about 35 mg of caffeine). Each caffeinated beverage was dichotomized (yes vs. no).

Other covariates

Alcohol consumption was defined as low (<1 alcoholic beverage a week), moderate (1 to 19 alcoholic beverages a week) and high to excessive consumption (>19 alcoholic beverages a week) (World Health Organization (WHO), 2004). Other covariates considered were: age (years), sex, cigarette smoking history (never, former, current) and participation in moderate or vigorous physical activity (no vs. yes); Body mass index (BMI) was calculated as weight (in kilograms) divided by height (in meters) squared. BMI thresholds were set according to previously defined WHO cutoff points (underweight, <18.5 kg/m²; normal, 18.5 to 24.9 kg/m²; overweight, 25.0 to 29.9 kg/m²; and obese, 30 kg/m²) (WHO, 1995).

Epworth sleepiness scale (ESS)

The Epworth sleepiness scale (ESS) was used to evaluate general level of daytime sleepiness (Johns, 1991) and the capability to stay alert and awake during crucial moments of the day (AASM, 2001). The ESS is a brief instrument that has been widely used globally among different study populations. It has 8 items capturing an individual's propensity to fall asleep during commonly encountered situations, each measured on a scale from 0 to 3. The scores for the eight questions are added together to obtain a single total score that ranges from 0 to 24. In adults, an ESS score ≥ 10 is taken to indicate increased daytime sleepiness (Johns, 1991). In this study we used the ESS ≥ 10 cutoff point to define daytime sleepiness.

The Horne and Ostberg morningness-eveningness questionnaire (MEQ)

Circadian preference was assessed using the Horne and Ostberg morningness-eveningness questionnaire (Horne et al., 1976). Used globally, the MEQ (Horne et al., 1976) is a 19-item questionnaire that identifies morningness-eveningness preference (Merikanto et al., 2013; Schneider et al., 2011). Circadian rhythm is an individual's endogenous sleep-wake state during a 24-h period, while circadian preference refers to a person's inclination to sleep and engage in activities during that same period (Barion and Zee, 2007; Horne et al., 1976; Schneider et al., 2011). The scores range from 16 to 86 and participants can be classified in five categories: definite and moderate evening (E)-type, neutral type and moderate and definite morning (M)-type. Higher values on MEQ indicate stronger morningness preference. In this study we used the following cut offs: (1) 16 to 41 for evening; (2) 42 to 58 for intermediate; (3) ≥ 59 for morning. In this study, we excluded intermediate types from analysis when comparing morningness and eveningness chronotypes.

Statistical analysis

We first examined the frequency distributions of socio-demographic and behavioral characteristics of the study participants.

Characteristics were summarized using means (\pm standard deviation) for continuous variables with normal distribution and counts and percentages for categorical variables. Mean (\pm standard deviation) of MEQ and ESS scores were calculated across socio-demographic and behavioral characteristics and the associations were tested using a one-way ANOVA for multi-level characteristics and a two-sample t-test for two-level characteristics. Multivariable linear regression models were also fitted to evaluate the associations. We also calculated the distribution of morningness-eveningness chronotype across demographic and behavioral characteristics and Chi-square tests were used to determine bivariate differences. Additionally, we calculated the distribution of daytime sleepiness and evening chronotype across energy drink consumption status. Multivariable logistic regression models were used to calculate odds ratios (ORs) and 95% confidence intervals (95% CIs) for the associations. All analysis were performed using SPSS Statistical Software for Windows (IBM SPSS, version 20, Chicago, IL, USA). All reported p-values are two-sided and deemed statistically significant at a 0.05 level.

RESULTS

Of the 3,000 college students who completed the survey and met participant guidelines, 66.9% of the students were females and the average reported age was 20.3 ± 1.3 years (Table 1). About two-thirds of students reported never drinking, while 32.1% admitted drinking 1 to 19 drinks per month; very few (1.7%) used alcohol excessively. A large majority of students have normal BMI (68.9%), few are overweight (10%) and 16.5% are underweight. Over three-fourths of study participants reported engaging in some type of physical activity. Table 2 shows the prevalence estimates of chronotype levels according to the Horne and Ostberg criteria. Thirteen percent of students were classified as evening types (12% in females and 15.1% in males) while 18.7% were classified as morning types (19.7% in females and 16.5% in males).

The prevalence of daytime sleepiness across age groups and sex is displayed in Figure 1. Daytime sleepiness (ESS ≥ 10) was present in 27.9% of the students (95% CI: 26.2 to 29.5%). Overall males in the 18 and the 22-and-over age groups appear to have higher prevalence of daytime sleepiness compared to females; 27.6% of the 18 year-old males experience daytime sleepiness compared to 21.2% of females; 35% of males 22-and-over experience daytime sleepiness compared to 27.4% of females. For the other age groups, the prevalence of daytime sleepiness is higher among females. Looking at the distribution of ESS total score across sex, the median ESS total score of females are higher than males.

As shown in Table 3, females (vs. males) and smokers (current and former vs. non-smokers) had a significantly lower MEQ score (p value < 0.001). There is also a significant association between MEQ score and alcohol consumption (p value < 0.001); we noted a trend lower MEQ scores with higher levels of alcohol consumption. Age, obesity status and participation in physical activity

Table 1. Characteristics of study sample.

Characteristic	N = 3,000	%
Age (Mean± SD)		20.3±1.3
Age (years)		
18	162	5.4
19	705	23.5
20	860	28.6
21	728	24.3
≥ 22	545	18.2
Sex		
Female	2,008	66.9
Male	992	33.1
Cigarette smoking status		
Never	2,739	91.3
Former	55	1.8
Current	206	6.9
Alcohol consumption		
<1 Drink/month	1,986	66.2
1-19 Drinks/month	962	32.1
≥ 20 Drinks/month	52	1.7
Body mass index (kg/m²)		
Underweight (<18.5)	495	16.5
Normal (18.5–24.9)	2,068	68.9
Overweight (25.0–29.9)	298	10
Obese (≥30.0)	139	4.6
Any physical activity		
No	669	22.4
Yes	2,319	77.6

Table 2. Prevalence estimates of morningness/eveningness chronotype

Parameter	MEQ score cut-off	All	Female	Male
		% (95% CI)		
Evening type (n=378)	≤41	13.0 (11.7-14.2)	12.0 (10.5-13.4)	15.1 (12.8-17.3)
Intermediate (n=1,984)	42-58	68.3 (66.6-69.9)	68.2 (66.1-70.3)	68.4 (65.4-71.3)
Morning type (n=544)	≥59	18.7 (17.3-20.1)	19.7 (18.0-21.5)	16.5 (14.1-18.9)

were not found to be statistically significantly associated with MEQ scores. The multivariable linear regression model with all the demographic and lifestyle characteristics gave similar results, except that sex was no longer found to be significant (Table 1).

As shown in Table 4, females (vs. males), alcohol consumers (vs. non-consumers), physically active respondents

(vs. non-physically active respondents) and non-obese individuals (vs. obese individuals) had a significantly higher ESS score (p value < 0.037). Age and smoking status was not found to be significantly associated with ESS score. The results from the multivariable linear regression model were not appreciably different from the univariate analyses (Table 2).

Table 3. Morningness/eveningnessquestionnaire (MEQ) scores by demographic and lifestyle characteristics.

Parameter	MEQ score		p-value
	Mean	SD	
Age (years)			
18	50.70	8.28	0.241
19	51.66	8.02	
20	50.87	8.09	
21	50.75	8.06	
≥ 22	50.97	8.44	
Sex			
Female	51.45	8.07	<0.001
Male	50.19	8.23	
Cigarette smoking status			
Never	51.40	8.00	<0.001
Former	47.56	8.97	
Current	47.02	8.56	
Alcohol consumption			
<1 Drink/month	52.14	8.07	<0.001
1-19 Drinks/month	49.04	7.75	
≥ 20 Drinks/month	45.31	8.76	
Obesity (≥30.0 kg/m²)			
No	51.07	8.15	0.334
Yes	50.38	7.99	
Any physical activity			
No	51.20	8.23	0.614
Yes	51.01	8.11	

Table 4. Epworth sleepiness scale (ESS) scores by demographic and lifestyle characteristics.

Parameter	EES score		p-value
	Mean	SD	
Age (years)			
18	7.44	2.84	0.635
19	7.56	3.35	
20	7.54	3.33	
21	7.57	3.47	
≥ 22	7.79	3.52	
Sex			
Female	7.68	3.32	0.037
Male	7.40	3.50	
Cigarette smoking status			
Never	7.54	3.36	0.059
Former	8.11	3.53	
Current	8.05	3.56	
Alcohol consumption			
<1 drink/month	7.46	3.39	0.009
1-19 drinks/month	7.84	3.32	
≥ 20 drinks/month	8.12	3.65	
Obesity (≥30.0 kg/m²)			
No	7.62	3.39	0.165
Yes	7.38	3.31	
Any physical activity			
No	7.18	3.45	0.001
Yes	7.70	3.34	

Table 5 summarizes the logistic regression results. Consumers of any stimulant beverage had 2.68-folds higher odds of being evening chronotypes compared to those who abstained from consuming stimulant beverages (OR 2.68; 95% CI 2.01 to 3.58), after adjusting for age, sex, smoking, BMI and physical activity. When considering specific types of beverages, the odds of being evening chronotype were between 1.95- and 3.5-fold higher among users compared to non-users. Compared to those who consumed less than one stimulant beverages per week, those who consumed two per week have 2.65-folds higher odds of being evening chronotypes (OR 2.65; 95% CI 1.81 to 3.90) while those who consumed three or more have 3.65-folds higher odds of being evening chronotypes (OR 3.65; 95% CI 2.58 to 5.16). However, consuming one stimulant per week was not statistically significantly associated with the odds of being evening chronotypes (OR 1.41; 95% CI .87 to 2.29). Con-sumers of stimulant beverages had a 22%

higher odds of experiencing daytime sleepiness compared to non-consumers (OR 1.22; 95% CI 1.03 to 1.44), after adjusting for demographic and lifestyle characteristics. When considering specific types of beverages, the odds of experiencing daytime sleepiness were between 1.21- and 1.72-fold higher among users compared to non-users; exceptions included coffee drinkers or consumers of other types of energy drinks, for whom no statistically significant association were found. Those who consumed three or more stimulants per week had 37% higher odds of experiencing daytime sleepiness compared to those who do not use stimulants (OR 1.37; 95% CI 1.13 to 1.67).

DISCUSSION

To the best of our knowledge, this is the first study to examine the prevalence of daytime sleepiness and evening




Figure 1. Prevalence of daytime sleepiness by age and sex.

Table 5. Evening chronotype and daytime sleepiness in relation to stimulant use.

Parameter	Evening Chronotype				Daytime Sleepiness			
	Yes (N=378)	No (N=544)	Unadjusted OR (95% CI)	Adjusted* OR (95% CI)	Yes (N=831)	No (N=2,152)	Unadjusted OR (95% CI)	Adjusted* OR (95% CI)
Any stimulant beverages	%	%			%	%		
No	29.6	54.6	1.00 (Reference)	1.00 (Reference)	38.6	43.7	1.00 (Reference)	1.00 (Reference)
Yes	70.4	45.4	2.86 (2.16-3.77)	2.68 (2.01,3.58)	61.4	56.3	1.24 (1.05-1.45)	1.22 (1.03-1.44)
Type of beverage								
Coffee	32.5	19.1	2.04 (1.51-2.76)	1.95 (1.42-2.67)	26.2	25.0	1.07 (0.89-1.28)	1.07 (0.89-1.29)
Tea	57.4	37.1	2.28 (1.75-2.98)	2.31 (1.75-3.05)	51.3	46.1	1.23 (1.05-1.45)	1.21 (1.03-1.43)
Coke/Pepsi with sugar	57.1	31.4	2.91 (2.22-3.82)	2.70 (2.03-3.58)	49.2	43.1	1.28 (1.09-1.50)	1.26 (1.07-1.48)
Coke/Pepsi sugar free	19.8	8.3	2.75 (1.847-4.08)	2.66 (1.77-4.00)	14.7	11.1	1.38 (1.09-1.74)	1.39 (1.10-1.76)
M 100/M 150	11.6	2.9	4.35 (2.41-7.83)	3.50 (1.90-6.44)	7.9	5.3	1.56 (1.14-2.13)	1.52 (1.10-2.11)
Red Bull	4.8	1.7	2.97 (1.32-6.69)	2.39 (1.02-5.58)	3.9	2.2	1.76 (1.11-2.77)	1.72 (1.08-2.75)
Other Energy Drinks ^{**}	3.4	0.9	3.84 (1.36-10.86)	2.90 (0.98-8.62)	2.8	1.6	1.72 (1.01-2.93)	1.67 (0.97-2.88)
Number of different types of stimulants/week								
0	29.6	54.6	1.00 (Reference)	1.00 (Reference)	38.6	43.7	1.00 (Reference)	1.00 (Reference)
1	8.5	11.0	1.41 (0.87-2.29)	1.21 (0.73-2.00)	8.8	9.5	1.04 (0.78-1.40)	1.03 (0.76-1.39)
2	22.2	15.3	2.68 (1.85-3.90)	2.65 (1.81-3.90)	18.5	18.8	1.12 (0.89-1.40)	1.10 (0.87-1.38)
≥3	39.7	19.1	3.83 (2.75-5.327)	3.65 (2.58-5.16)	34.1	27.9	1.38 (1.14-1.67)	1.37 (1.13-1.67)

*Adjusted for age, sex, smoking, body mass index, and physical activity; ** Other energy drinks includes the following: CarabaoDaeng, Lipovitan-D or Lipo, Wrangyer and Shark.

evening chronotype in relation to caffeine use in a Southeast Asian population. Approximately 28% experienced daytime sleepiness while 13% of our student cohort reported to be evening types. Overall, students with daytime sleepiness were more likely to be cigarette smokers, alcohol drinkers, obese, physically active and stimulant beverage consumers. The odds of being evening types were elevated among cigarette smokers, alcohol consumers and caffeinated stimulant users.

Our results are in accordance with previous reports indicating prevalence of daytime sleepiness and eveningness chronotype and extend this literature to assess their associations with consumption of stimulant beverages in Thai young adults (Adan and Natale, 2002; Besoluk et al., 2011; Chung and Cheung, 2008; Hsu et al., 2012; Lund et al., 2010; Schneider et al., 2011; Wu et al., 2012; Yang et al., 2003). For instance, our result showing that 13% of Thai college students were evening types is consistent with reports from Chung and Cheung (2008) who reported a 13.1% prevalence of evening chronotype among Chinese students. Additionally, our findings showing a 27.9% daytime sleepiness are similar to those of Wu et al. (2012) who reported a 22.2% prevalence of daytime sleepiness. Other investigators, however, have reported higher prevalence estimates (Pirrallo et al., 2012). On balance, the results of our study and those of others emphasize the growing problem of daytime sleepiness and eveningness chronotype among college students.

In our study of Thai college students, consuming any stimulant beverage was strongly associated with evening chronotype and daytime sleepiness. Prior studies have found similar results in other university settings (Giannotti et al., 2002; James et al., 2011; Roehrs and Roth 2008; Snel and Lorist, 2011; Taylor et al., 2011). College marks the beginning of new and stressful changes: increased academic workload, busier social lives and later bed times (Lund et al., 2010; Malinauskas et al., 2007). Several investigators determined that academic stress interferes with sleeping schedule, as students may stay up late into the night to study (Chung et al., 2008; Lund et al., 2010). Given the high prevalence of poor sleep quality, short sleep duration and high rates of energy drink consumption among Thai college students, (Steptoe et al., 2006; Zenith International, 2007) it may be fitting to note that these undergraduates may sacrifice sleep for academic purposes and social commitments and drink caffeinated stimulants to remain alert (Ishak et al., 2012; Malinauskas et al., 2007; Roehrs and Roth 2008). As they stay up later to finish their work, they can become sleep deprived due to the morning-oriented schedules of college and thus experience daytime sleepiness (Kushida, 2006; Wittmann et al., 2010).

Several investigators have noted that evening types and daytime sleepers are more likely to have poor health (AASM, 2001; Kanerva et al., 2012; Killgore et al., 2008; Nakade et al., 2009; Schneider et al., 2011; Slater et al.,

2013; Stroe et al., 2010; Wittmann et al., 2010). Our findings suggest that caffeine consumers were more likely to experience daytime sleepiness and be evening types. As a result, caffeine could possibly be linked to other poor health behaviors. Future research must evaluate the extent to which caffeinated beverages and energy drinks are associated with poor health traits.

The association between caffeinated drinks and circadian disruption and daytime sleepiness can be explained by the biological mechanism of melatonin suppression and adenosine blockage. During waking hours, light is known to suppress melatonin production (Barion et al., 2007; Gooley et al., 2011). Given the demanding college workload and social commitments, students may stay up later into the night to study, which requires the use of lighting and encourages the consumption of caffeinated beverages to increase alertness. This increased light exposure leads to melatonin suppression, in which light exposure at night shifts one's chronotype towards eveningness (Barion et al., 2007; Shanahan et al., 1999). Another mechanism that can explain sleep disruption is caffeine's role in adenosine blockage. Within the basal forebrain, adenosine, an endogenous biochemical compound, regulates sleep by inhibiting the cholinergic neurons that create arousal (Basheer et al., 2004; Ribeiro and Sebastiao, 2010; Roehrs and Roth 2008). As caffeine acts as an adenosine blocker, it negates the effects of adenosine to induce sleep (Roehrs and Roth 2008). Consequently, consuming caffeine at night prolongs wakefulness and decreases sleep duration, resulting in daytime sleepiness (James et al., 2011; Kushida, 2006; Roehrs and Roth 2008).

The results from our study should be interpreted in the context of some limitations. First, our study could be subjected to volunteer bias, because the data were collected from willing participants instead of a random sample. Second, the temporal relationship between lifestyle characteristics and sleep disorders cannot be delineated due to the cross-sectional study design. It is possible that daytime sleepiness and eveningness chronotype might have led to increased consumption of energy drinks and other caffeinated beverages. Prospective studies are required to confirm and expand upon our observations. Third, there may be lifestyle and dietary traits that are heterogeneous within daytime sleepers and evening types, which could affect the strength of the associations between sleep disorders and lifestyle traits. Fourth, our study population was exclusively comprised of participants who were attending college, thus conclusions cannot be other broader populations. Lastly, we did not have information concerning frequency, timing and dose of energy drink consumption in the present study. As a result, it is possible that the binary grouping of energy drink consumption may have attenuated the magnitude of the association toward the null.

Conclusion

In summary, our findings underline the growing problem of energy drinks and their impact on both circadian preference and daytime sleepiness among Thai college students. Despite academics being the frequent reason given for energy drink consumption, evening types or daytime sleepers do not exhibit higher academic performance compared to their counterparts (Besoluk et al., 2011; Chung et al., 2008; Giannotti et al., 2002; James et al., 2011; Taylor et al., 2011).

RECOMMENDATION

There is a large body of evidence that daytime sleepers and evening types are strongly associated with numerous health and social problems (AASM, 2001; Kanerva et al., 2012; Killgore et al., 2008; Nakade et al., 2009; Schneider et al., 2011; Slater et al., 2013). From a public health promotion and disease prevention standpoint, these findings suggest an obvious need for effective educational and prevention programs targeted toward improving sleep hygiene and reducing consumption of energy drinks among young adults. Future research that evaluates the impact of caffeinated beverages on various sleep disorders and those that assess the effectiveness of health and wellness programs among young adults, are needed.

ACKNOWLEDGEMENT

This research was completed while Mr. Jason Tran was a research training fellow with the Harvard School of Public Health Multidisciplinary Health International Research Training (HSPH MHIRT) Program. The HSPH MHIRT Program is supported by an award from the National Institute for Minority Health and Health Disparities (T37-MD000149).

Conflict of interest

The authors have no competing interests to declare.

REFERENCES

- AASM (2001). International Classification of Sleep Disorders, revised: Diagnostic and coding manual. Chicago, Illinois: American Academy of Sleep Medicine.
- Adan A, Natale V (2002). Gender differences in morningness-eveningness preference. *Chronobiol Int* 19(4):709-720.
- Arria AM, Caldeira KM, Kasperski SJ, O'Grady KE, Vincent KB, Griffiths RR, Wish ED (2010). Increased alcohol consumption, nonmedical prescription drug use, and illicit drug use are associated with energy drink consumption among college students. *J. Addict. Med.* 4(2):74-80.
- Barion A, Zee PC (2007). A clinical approach to circadian rhythm sleep disorders. *Sleep Med.* 8(6):566-577.
- Basheer R, Strecker RE, Thakkar MM, McCarley RW (2004). Adenosine and sleep-wake regulation. *Prog. Neurobiol.* 73(6):379-396.
- Besoluk S, Onder I, Deveci I (2011). Morningness-eveningness preferences and academic achievement of university students. *Chronobiol Int.* 28(2):118-125.
- Buxton C, Hagan JE (2012). A survey of energy drinks consumption practices among student -athletes in Ghana: lessons for developing health education intervention programmes. *J. Int. Soc. Sports Nutr.* 9(1):9.
- Cannon ME, Cooke CT, McCarthy JS (2001). Caffeine-induced cardiac arrhythmia: an unrecognized danger of healthfood products. *Med. J. Aust.* 174(10):520-521.
- Cheng SH, Shih CC, Lee IH, Hou YW, Chen KC, Chen KT, Yang YK, Yang YC (2012). A study on the sleep quality of incoming university students. *Psychiatry Res.* 197(3):270-274.
- Chung KF, Cheung MM (2008). Sleep-wake patterns and sleep disturbance among Hong Kong Chinese adolescents. *Sleep* 31(2):185-194.
- Doi Y, Inoue Y, Minowa M, Uchiyama M, Okawa M (2003). Periodic leg movements during sleep in Japanese community-dwelling adults based on the assessments of their bed partners. *J. Epidemiol.* 13(5):259-265.
- Giannotti F, Cortesi F, Sebastiani T, Ottaviano S (2002). Circadian preference, sleep and daytime behaviour in adolescence. *J. Sleep Res.* 11(3):191-199.
- Gooley JJ, Chamberlain K, Smith KA, Khalsa SB, Rajaratnam SM, Van Reen E, Zeitzer JM, Czeisler CA, Lockley SW (2011). Exposure to room light before bedtime suppresses melatonin onset and shortens melatonin duration in humans. *J. Clin. Endocrinol. Metab.* 96(3):E463-472.
- Horne JA, Ostberg O (1976). A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Chronobiol. Int.* 4(2):97-110.
- Hsu CY, Gau SS, Shang CY, Chiu YN, Lee MB (2012). Associations between chronotypes, psychopathology, and personality among incoming college students. *Chronobiol Int.* 29(4):491-501.
- Ishak WW, Ugochukwu C, Bagot K, Khalili D, Zaky C (2012). Energy drinks: psychological effects and impact on well-being and quality of life-a literature review. *Innov. Clin. Neurosci.* 9(1):25-34.
- James JE, Kristjansson AL, Sigfusdottir ID (2011). Adolescent substance use, sleep, and academic achievement: evidence of harm due to caffeine. *J. Adolesc.* 34(4):665-673.
- Johns MW (1991). A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 14(6):540-545.
- Kanerva N, Kronholm E, Partonen T, Ovaskainen ML, Kaartinen NE, Kontinen H, Broms U, Männistö S (2012). Tendency toward eveningness is associated with unhealthy dietary habits. *Chronobiol. Int.* 29(7):920-927.
- Killgore WD, Kahn-Greene ET, Lipizzi EL, Newman RA, Kamimori GH, Balkin TJ (2008). Sleep deprivation reduces perceived emotional intelligence and constructive thinking skills. *Sleep Med.* 9(5):517-526.
- Kushida CA (2006). Countermeasures for sleep loss and deprivation. *Curr. Treat Options Neurol.* 8(5):361-366.
- Lohsoonthorn V, Khidir H, Casillas G, Lertmaharit S, Tadesse MG, Pensuksan WC, Rattananupong T, Gelaye B, Williams MA (2012). Sleep quality and sleep patterns in relation to consumption of energy drinks, caffeinated beverages, and other stimulants among Thai college students. *Sleep Breath* 17:1017-28.
- Lucassen EA, Zhao X, Rother KI, Mattingly MS, Courville AB, de Jonge L, Csako G, Cizza G; Sleep Extension Study Group (2013). Evening chronotype is associated with changes in eating behavior, more sleep apnea, and increased stress hormones in short sleeping obese individuals. *PLoS One* 8(3):e56519.
- Lund HG, Reider BD, Whiting AB, Prichard JR (2010) Sleep patterns and predictors of disturbed sleep in a large population of college students. *J. Adolesc. Health* 46(2):124-132.
- Luyster FS, Strollo PJ Jr, Zee PC, Walsh JK; Boards of Directors of the American Academy of Sleep Medicine and the Sleep Research Society (2012). Sleep: a health imperative. *Sleep* 35(6):727-734.
- Mak KK, Ho SY, Thomas GN, Lo WS, Cheuk DK, Lai YK, Lam TH (2010). Smoking and sleep disorders in Chinese adolescents. *Sleep*

- Med. 11(3):268-273.
- Malinauskas BM, Aeby VG, Overton RF, Carpenter-Aeby T, Barber-Heidal K (2007). A survey of energy drink consumption patterns among college students. *Nutr. J.* 6:35.
- Merikanto I, Lahti T, Kronholm E, Peltonen M, Laatikainen T, Vartiainen E, Salomaa V, Partonen T (2013). Evening types are prone to depression. *Chronobiol. Int.* 30(5):719-725.
- Millman RP (2005). Excessive sleepiness in adolescents and young adults: causes, consequences, and treatment strategies. *Pediatrics* 115(6):1774-1786.
- Nakade M, Takeuchi H, Kurotani M, Harada T (2009). Effects of meal habits and alcohol/cigarette consumption on morningness-eveningness preference and sleep habits by Japanese female students aged 18-29. *J Physiol. Anthropol.* 28(2):83-90.
- Oginska H, Pokorski J (2006). Fatigue and mood correlates of sleep length in three age-social groups: School children, students, and employees. *Chronobiol. Int.* 23(6):1317-1328.
- Pirralo RG, Loomis CC, Levine R, Woodson BT (2012). The prevalence of sleep problems in emergency medical technicians. *Sleep Breath* 16(1):149-162.
- Pomeranz JL, Munsell CR, Harris JL (2013). Energy drinks: an emerging public health hazard for youth. *J. Public Health Policy* 34(2):254-271.
- Ribeiro JA, Sebastiao AM (2010). Caffeine and adenosine. *J. Alzheimers Dis.* 20 Suppl 1:S3-15.
- Roehrs T, Roth T (2008). Caffeine: sleep and daytime sleepiness. *Sleep Med. Rev.* 12(2):153-162.
- Schneider ML, Vasconcellos DC, Dantas G, Levandovski R, Caumo W, Allebrandt KV, Hidalgo MP (2011). Morningness-eveningness, use of stimulants, and minor psychiatric disorders among undergraduate students. *Int. J. Psychol.* 46(1):18-23.
- Seifert SM, Schaechter JL, Hershorin ER, Lipshultz SE (2011). Health effects of energy drinks on children, adolescents, and young adults. *Pediatrics* 127(3):511-528.
- Shanahan TL, Kronauer RE, Duffy JF, Williams GH, Czeisler CA (1999). Melatonin rhythm observed throughout a three-cycle bright-light stimulus designed to reset the human circadian pacemaker. *J. Biol. Rhythms* 14(3):237-253.
- Slater G, Pengo MF, Kosky C, Steier J (2013). Obesity as an independent predictor of subjective excessive daytime sleepiness. *Respir. Med.* 107(2):305-309.
- Snel J, Lorist MM (2011). Effects of caffeine on sleep and cognition. *Prog. Brain Res.* 190:105-117.
- Steptoe A, Peacey V, Wardle J (2006). Sleep duration and health in young adults. *Arch. Intern. Med.* 166(16):1689-1692.
- Stroe AF, Roth T, Jefferson C, Hudgel DW, Roehrs T, Moss K, Drake CL (2010). Comparative levels of excessive daytime sleepiness in common medical disorders. *Sleep Med.* 11(9):890-896.
- Taillard J, Philip P, Bioulac B (1999). Morningness/eveningness and the need for sleep. *J. Sleep Res.* 8(4):291-295.
- Taylor DJ, Clay KC, Bramoweth AD, Sethi K, Roane BM (2011). Circadian phase preference in college students: relationships with psychological functioning and academics. *Chronobiol. Int.* 28(6):541-547.
- Tsai LL, Li SP (2004). Sleep patterns in college students: gender and grade differences. *J. Psychosom. Res.* 56(2):231-237.
- WHO (1995). Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. World Health Organization, Geneva.
- WHO (2004). Global status report on alcohol. World Health Organization, Department of Mental Health and Substance Abuse, Geneva.
- Wittmann M, Paulus M, Roenneberg T (2010). Decreased psychological well-being in late 'chronotypes' is mediated by smoking and alcohol consumption. *Subst Use Misuse* 45(1-2):15-30.
- Wu S, Wang R, Ma X, Zhao Y, Yan X, He J (2012). Excessive daytime sleepiness assessed by the Epworth Sleepiness Scale and its association with health related quality of life: a population-based study in China. *BMC Public Health*, 12:849.
- Yang CM, Wu CH, Hsieh MH, Liu MH, Lu FH (2003). Coping with sleep disturbances among young adults: a survey of first-year college students in Taiwan. *Behav. Med.* 29(3):133-138.
- Zenith International (2007). US overtakes Thailand as world leader in energy drinks. [June 1, 2013]. http://www.zenithinternational.com/news/press_release_detail.asp?id=206.